Application Note FOB Turbilatex, Chemwell-T, Awareness (AN-Fb-ChemWell®-T.EN rev 2019.05.27)

For in vitro diagnostic device ENGLISH





General Information

Intended use:

FOB Turbilatex is a latex turbidimetric assay for the quantitative detection of human haemoglobin (hHb) in human stool samples.

This assay is simple and widely applicable. Test results aid in a presumptive diagnosis of faecal occult blood (gastrointestinal bleeding).

For professional in vitro diagnostic use only.

FOB Turbilatex can be performed on every open chemistry analyser. Please follow the subsequent instructions in order to assure performance characteristics as describes in the instructions for use. This instruction has been validates by CerTest BIOTEC S.L Laboratories.

Additionally please read the "Instructions for use" for instructions on operating and programming user defined test.

Reagents:

Materials provided by CerTest BIOTEC S.L.:

Reagents	Quantity	Cat. reference
Turbidimetric reagents (R1 & R2) 200 Det/kit	R1: 2 vials, 2x15 mL. R2: 1 vial, 1x6 mL.	TL-022FB01C TL-022FB02C
Auxiliary Reagents	Quantity	Cat. reference
Calibration kit	Calibrator: 6 vials, 6x1mL.	TL-022FB70, TL-022FB71, TL-022FB72, TL-022FB73, TL-022FB74, TL-022FB75
Controls kit	Control C1, 2 vials, 2x1 mL/vial. Control C2, 2 vials, 2x1 mL/vial.	TL-022FB08 TL-022FB09
Sample diluent kit	4 vials, 4x125 mL/vial	TL-022FB03E
Sample dilutions vials	1x2 mL/vial 1x2 mL/vial	MST-0005MF MST-0009F

Preparation of reagents:

R1 and R2 are ready to use.

Calibrators are ready to use.

Controls are ready to use.

Storage and stability

Kit components must be stored at temperature indicated on the label. Do not freeze.

Reagents are stable up to the expiration date printed on the label, always considering that reagent containers must be properly closed to avoid any contamination, must be kept away from the sunlight and conserved at temperature indicated on the label of each reagent.

Specimen:

Collect enough quantity of human stool samples. These samples should be collected in clean and dry containers (no preservatives or transport media). The samples can be stored completely dissolved in the refrigerator (2-8°C) for 3 days prior to testing. Homogenise stool samples as thoroughly as possible prior to preparation.

The sample dilution vial with diluted sample can be stored for 7 days in the refrigerator (2-8°C) prior to testing.

Use FOB Turbilatex stool collection tubes for sample collections described the instructions for use.

Assay procedure

Application parameter set up:

Specific analyzers settings for FOB Turbilatex must be programmed onto the analyzer, see below. For instructions, consult the ChemWell®-T (Awareness Technology Inc.) analyzer manual and instructions for use provided with the kits.

Loading of reagents:

Load reagents according to the ChemWell®-T (Awareness Technology Inc.) analyzer manual.

Calibration curve establishment:

A 6 points calibration curve can be established in ChemWell®-T (Awareness Technology Inc.) analyzer. For instructions consult analyzer manual.

Calibration stability:

Calibrate the system at least once a month is extremely recommended. Recalibrate the system when reagent lot is change or when the controls are out of the assigned range given in the control label and CoA.

QC controls:

FOB Turbilatex controls C1 and C2 must be assayed each day before running patient fecal sample extract to validate the calibration curve. The controls have assigned value ranges indicated on the label and certificate of analysis supplied. The control measurements must be within the indicated value range to obtain valid results for patient fecal extract. If the control values are out of range, follow next procedures:

1) Repeat QC control measurement, 2) Repeat calibration measurement.

Results

The results are evaluated automatically by the analyzer and presented in ng hHb/mL.

FOB Turbilatex, Chemwell-T, Awareness

(AN-Fb-ChemWell®-T.EN rev 2019.05.27)

For *in vitro* diagnostic device ENGLISH






Performance characteristics

The following results have been obtained during the validation of FOB Turbilatex on the ChemWell®-T (Awareness Technology Inc.) analyzer.

Linearity:

FOB Turbilatex on ChemWell®-T (Awareness Technology Inc.) analyzer using calibrator kit is linear in the calibration range of 0-1000 ng hHb/mL.



Measuring range:

FOB Turbilatex assay measuring range is 15-1000 ng hHb/mL on the ChemWell®-T (Awareness Technology Inc.) analyser. Samples higher concentrated than 1000 ng hHb/mL must be diluted for proper quantification by the user, using additional sample buffer.

Prozone effect

Using the reported parameters, no hook effect was observed up to 10 μ g hHb/mL. Samples with Haemoglobin concentration of 16 μ g hHb/mL give a typical positive result >1000 ng hHb/mL.

Detection limit

Limit of detection (LOD): 13 ng hHb/mL. The lower limit of detection of FOB Turbilatex was determined on 20 samples and 2 sample replicates as the mean value $+ 2 \cdot SD$.

Limit of quantification (LOQ): 15 ng hHb/mL. The lower limit of quantification is defined as the lowest actual amount of analysis that can be reliably detected; imprecision is < 20% as CV% on the ChemWell®-T (Awareness Technology Inc.) analyzer.

Precision

FOB Turbilatex was tested with three different controls levels.

	Low (20 ng/mL)	Medium (80 ng/mL)	High (250 ng/mL)
N	20	20	20
Mean (ng/mL)	22.0	84.2	254.2
SD (ng/mL)	2.1	7.1	9.3
CV (%)	9	8	4

Method comparison

Reults obtained with FOB Turbilatex on the ChemWell®-T (Awareness Technology Inc.) analyzer were compared with those obtained with EIKEN FOB Latex.

	Sensitivity	Specificity
FOB Turbilatex vs EIKEN FOB Latex	96%	>99%

Shipping damage

Please notify your distributor, it this product was received damaged.

Symbols key

IVD	For in vitro diagnostic use only	*	Keep dry
[]i	Consult instructions for use	1	Temperature limitation
REF	Catalogue number	LOT	Lot number
53	Use by	ш	Manufacturer
\sum_{n}	Contains sufficient for <n> test</n>	DIL	Sample diluent
誉	Keep out of the sunlight		

Manufacturer

CERTEST BIOTEC

Pol. Industrial Río Gállego II,Calle J, № 1, 50840, San Mateo de Gállego, Zaragoza (SPAIN) www.certest.es

NOTES

Please refer to the instruction for use for the detailed information about the test on the following:

Synthesis; Principle; Precautions; Reagents; Specimen collection; Interpretation of results.



ChemWell-T (Awareness) / Application parameters

Sample 20 R2 40 PR2 40 PR2 40 PR2 A0	ASSAY PARAMETERS	
Sample 20 R2 40 PR2 40 PR2 40 PR2 40 PR2 A0	Std. No	6
Reaction mode Endpoi Primary wavelength 505 n Secondary wavelength Nor Direction Increa: Reagent blank lecture 10 seconds after R2 addition Final lecture 300 seconds after blank lecture Reaction time 10 m Linear range 0-1000 ng/n CALIBRATION Calibration Method Polinomial 4th ord Calibration set 6 calibrato Blank Calibrator 1 (0 ng/m Calibrator 1 Calibrator 2 (50 ng/m Calibrator 3 Calibrator 3 (100 ng/m Calibrator 4 Calibrator 5 (500 ng/m Calibrator 5 Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	R1	150 μL
Reaction mode Endpoi Primary wavelength 505 m Secondary wavelength Nor Direction Increase Reagent blank lecture 10 seconds after R2 addition Final lecture 300 seconds after blank lectur Reaction time 10 m Linear range 0-1000 ng/m CALIBRATION Calibration Method Polinomial 4th ord Calibration set 6 calibrator 1 Calibrator 1 Calibrator 2 (50 ng/m Calibrator 2 Calibrator 3 (100 ng/m Calibrator 3 Calibrator 4 (250 ng/m Calibrator 5 (500 ng/m Calibrator 5 Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	Sample	20 μL
Reaction mode Primary wavelength Scondary wavelength Secondary wavelength Nor Direction Reagent blank lecture Final lecture Reaction time Linear range CALIBRATION Calibration Method Polinomial 4 th ord Calibrator set Blank Calibrator 1 Calibrator 2 Calibrator 2 Calibrator 3 Calibrator 3 Calibrator 4 Calibrator 5 Calibrator 6 (1000 ng/m CSTEPS Addition Sample	R2	40 μL
Primary wavelength Secondary wavelength Nor Direction Reagent blank lecture Final lecture Reaction time Linear range O-1000 ng/r CALIBRATION Calibration Method Polinomial 4 th ord Calibrator 1 Calibrator 1 Calibrator 2 Calibrator 3 Calibrator 3 Calibrator 4 Calibrator 5 STEPS Addition R1 Addition Sample	Others	NA
Secondary wavelength Direction Increase Reagent blank lecture Final lecture Reaction time Linear range CALIBRATION Calibration Method Polinomial 4th ord Calibrator 1 Calibrator 2 Calibrator 2 Calibrator 3 Calibrator 3 Calibrator 4 Calibrator 5 Calibrator 5 Calibrator R1 Addition R1 Addition Sample	Reaction mode	Endpoint
Direction Increase Reagent blank lecture 10 seconds after R2 addition Final lecture 300 seconds after blank lecture Reaction time 10 m Linear range 0-1000 ng/m CALIBRATION Calibration Method Polinomial 4th ord Calibration set 6 calibrator 1 (0 ng/m Calibrator 1 Calibrator 2 (50 ng/m Calibrator 2 Calibrator 3 (250 ng/m Calibrator 4 (250 ng/m Calibrator 5 (500 ng/m Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	Primary wavelength	505 nm
Reagent blank lecture Final lecture Reaction time Linear range CALIBRATION Calibration Method Polinomial 4 th ord Calibrator set Blank Calibrator 1 Calibrator 2 Calibrator 2 Calibrator 3 Calibrator 3 Calibrator 4 Calibrator 5 STEPS Addition R1 Addition Sample	Secondary wavelength	None
Final lecture Reaction time Linear range CALIBRATION Calibration Method Calibration set Blank Calibrator 1 Calibrator 2 Calibrator 2 Calibrator 3 Calibrator 3 Calibrator 4 Calibrator 5 STEPS Addition R1 Addition Sample	Direction	Increase
Reaction time Linear range O-1000 ng/r CALIBRATION Calibration Method Polinomial 4 th ord Calibration set Galibrator 1 Calibrator 1 Calibrator 2 Calibrator 2 Calibrator 3 Calibrator 3 Calibrator 4 Calibrator 4 Calibrator 5 Calibrator 5 Calibrator 6 (1000 ng/m Calibrator R1 Addition Sample	Reagent blank lecture	10 seconds after R2 addition
Linear range CALIBRATION Calibration Method Calibration set Blank Calibrator 1 (0 ng/m Calibrator 2 (50 ng/m Calibrator 3 (100 ng/m Calibrator 4 (250 ng/m Calibrator 5 (500 ng/m Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	Final lecture	300 seconds after blank lecture
CALIBRATION Calibration Method Calibration set Blank Calibrator 1 (0 ng/m Calibrator 2 (50 ng/m Calibrator 3 (100 ng/m Calibrator 4 (250 ng/m Calibrator 5 (500 ng/m STEPS Addition R1 Addition Sample	Reaction time	10 min
Calibration MethodPolinomial 4th ordCalibration set6 calibratorBlankCalibrator 1 (0 ng/mCalibrator 1Calibrator 2 (50 ng/mCalibrator 2Calibrator 3 (100 ng/mCalibrator 3Calibrator 4 (250 ng/mCalibrator 4Calibrator 5 (500 ng/mCalibrator 5Calibrator 6 (1000 ng/mSTEPSAddition R1Addition Sample	Linear range	0-1000 ng/ml
Calibration set Blank Calibrator 1 (0 ng/m Calibrator 2 (50 ng/m Calibrator 2 Calibrator 3 (100 ng/m Calibrator 3 Calibrator 4 Calibrator 4 Calibrator 5 Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	CALIBRATION	
Blank Calibrator 1 (0 ng/m Calibrator 2 (50 ng/m Calibrator 2 (50 ng/m Calibrator 3 (100 ng/m Calibrator 3 Calibrator 4 (250 ng/m Calibrator 4 Calibrator 5 (500 ng/m Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	Calibration Method	Polinomial 4 th order
Calibrator 1 Calibrator 2 (50 ng/m Calibrator 2 Calibrator 3 (100 ng/m Calibrator 3 Calibrator 4 (250 ng/m Calibrator 4 Calibrator 5 Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample		6 calibrators
Calibrator 2 Calibrator 3 (100 ng/m Calibrator 3 Calibrator 4 (250 ng/m Calibrator 5 Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	Blank	Calibrator 1 (0 ng/ml)
Calibrator 3 Calibrator 4 (250 ng/m Calibrator 4 Calibrator 5 (500 ng/m Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample	Calibrator 1	Calibrator 2 (50 ng/ml)
Calibrator 4 Calibrator 5 (500 ng/m Calibrator 5 (200 ng/m STEPS Addition R1 Addition Sample		Calibrator 3 (100 ng/ml)
Calibrator 5 Calibrator 6 (1000 ng/m STEPS Addition R1 Addition Sample		Calibrator 4 (250 ng/ml)
STEPS Addition R1 Addition Sample	Calibrator 4	Calibrator 5 (500 ng/ml)
Addition R1 Addition Sample	Calibrator 5	Calibrator 6 (1000 ng/ml)
Addition Sample	STEPS	
·	Addition R1	
	Addition Sample	
Incubation 5 m	Incubation	5 min
Addition R2	Addition R2	
Blank Lecture	Blank Lecture	
Incubation 5 m	Incubation	5 min
Final lecture	Final lecture	

Application Note


For help, press F1

FOB Turbilatex, Chemwell-T, Awareness

(AN-Fb-ChemWell®-T.EN rev 2019.05.27)

For *in vitro* diagnostic device ENGLISH





NUM